



## Aquatic Invasive Species (AIS) Control Plan: New Zealand Mudsnail

This control plan is a living document and will be updated, as needed, to reflect the status of the species within Pennsylvania.

### Natural History

Description: New Zealand Mudsnails (*Potamopyrgus antipodarum*) are diminutive freshwater gastropods, typically attaining adult sizes of only several millimeters (about 1/4<sup>th</sup> of an inch) in length.

### Taxonomy

Common name: **New Zealand Mudsnail**  
Family: **Tateidae**  
Species: *Potamopyrgus antipodarum*  
Integrated Taxonomic Information System (ITIS) Serial Number **205006**

Morphology: New Zealand Mudsnails (hereafter in this document abbreviated to “NZM”) are small freshwater snails which typically attain maximum lengths of only 4-6 mm within invaded portions of North America (Figure 1) but can attain larger sizes in their native range (Levri et al. 2007). NZM contain a “corkscrew” shaped spiraled shell that coils to the right and typically contains 5-8 whorls (Benson et al. 2023). NZM contain a hard operculum which can close to cover the opening in the shell. Shells are typically various shades of grey or brown. Populations within the Great Lakes contain different shell morphologies, such as a keel in the middle of each whorl, which is not typical of populations elsewhere (Levri

et al. 2007). Some populations also have small spines on the shell for defense from predators (Benson et al. 2023). NZM can be easily confused with other snail species and so experts are typically required to confirm identification.



**Figure 1.** Top: Magnified image of New Zealand Mudsnail. Source: USGS. Bottom: New Zealand Mudsnail shells with penny for scale. Source: Tim Throne, Trout Unlimited.

Origin: NZM are native to freshwater environments in New Zealand and several surrounding islands. NZM have spread to numerous localities in Australia, Europe, and North America. In the United States, NZM were first reported from near the Snake River, Idaho, in 1987 (Benson et al. 2023).



**Food Preferences:** NZM feed primarily on detritus, periphyton, and organic particles within sediment (Bilka and Levri 2013; Benson et al. 2023).

**Reproduction:** In their native range, NZM females primarily reproduce asexually via parthenogenesis, although sexual reproduction is known to occur. Invasive populations of NZM within the United States are thought to be exclusively clonal colonies of asexual females (Benson et al. 2023).

**Notable Characteristics:** NZM can achieve high densities within invaded waters (Figure 2); with some extreme estimates suggesting densities of over 299,000 snails per m<sup>2</sup> (Kerans et al. 2005).



**Figure 2.** Rock infested with New Zealand Mudsails from Codorus Creek, Pennsylvania. Credit: Tom Feneniz (Trout Unlimited).

**Historic Vectors:** NZM are thought to have spread from New Zealand to other parts of the world inadvertently through the global shipping trade. It is speculated that NZM were introduced to the Great Lakes on ships from Europe, and in water with game fish shipped from infested waters and stocked in western North America (Benson et al. 2023).


**Current Pathways/Vectors:** On smaller geographic scales (i.e., transport between watersheds), NZM are typically unintentionally introduced as “hitchhikers” on fishing, boating, and other aquatic recreational gear (Benson et al. 2023). NZM can close their operculum and can resist desiccation for up to several days outside of water (Richards et al. 2004). In Pennsylvania, the epicenter of most NZM infestations are located at popular trout fishing locations, suggesting waders/fishing gear as a major vector of transport (Hartzell and Macelko, in press). NZM may also be transported inadvertently via fish stocking and may even survive passage through the digestive system of some fish such as trout (Vinson and Baker 2008). NZM may also disperse on aquatic plants that are transported by humans or on aquatic vegetation moved during high water events (Benson et al. 2023).

**Preferred Habitat:** NZM are tolerant of a broad range of freshwater habitat characteristics and can even occur in brackish waters (Benson et al. 2023). They may be collected in deeper waters of lakes (Levri et al. 2007). In general, they prefer lentic or slow flowing waters with high nutrient levels, but are tolerant of fast flows, sedimentation, and pollution (Benson et al. 2023). In Pennsylvania, they appear to prefer waters with high conductivity (> 200  $\mu\text{S}/\text{cm}^2$ ) and basic (>7) pH levels, likely due to a greater availability of calcium ions required for shell growth at these conditions (Levri et al. 2020). Anecdotally, many of the populations of this species with high densities in Pennsylvania appear to be in coldwater streams with a limestone influence.



## Distribution and Status

**Distribution:** NZM have been reported in at least 21 US States, with populations primarily concentrated in the western US and in the vicinity of the Great Lakes (Benson et al. 2023; Figure 3).




**Figure 3.** Distribution of New Zealand Mudsnails within the continental United States. Source: USGS.

In Pennsylvania, NZM have been reported from at least 17 counties (Figure 4). NZM were first documented within Pennsylvania in 2005 from Lake Erie in the vicinity of Presque Isle State Park (Levri et al. 2007). In 2010, NZM were discovered in Spring Creek, Centre County (Levri et al. 2020). Since 2018, NZM have been documented in numerous streams and several rivers, many of which are popular trout waters within the Lake Erie, Susquehanna, Delaware, Allegheny/Ohio, and Potomac River basins in Pennsylvania (Benson et al. 2023; Hartzell and Frederick 2023; Hartzell and Macelko, in press). Complete information on the distribution of NZM in Pennsylvania, including specific waters, can be found at the USGS nonindigenous aquatic species webpage:

<https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=1008>

**Pennsylvania Legal Status:** As of November 2023, NZM are not legally regulated in Pennsylvania.



**Figure 4.** County-level distribution of New Zealand Mudsnails in Pennsylvania (November 2023).

## Threats

**Ecological:** Due to their potential to attain high densities within invaded waters, NZM are competitors with native gastropods and other periphyton-grazing macroinvertebrates such as caddisfly larvae and mayfly nymphs (Karens et al. 2010; Krist and Charles 2012; Larson and Black 2016) and NZM infestations may ultimately result in reductions of macroinvertebrate abundance and biodiversity (Karens et al. 2005; 2010), however, in some areas this impact may be minor (Cada 2004; Karens et al. 2005). Few studies appear to have been done evaluating the potential impacts of NZM on higher trophic levels (i.e., fish). However, trout will readily consume NZM in both laboratory and natural settings and will lose weight due to the poor nutritional content of NZM as compared to native macroinvertebrates (Vinson and Baker 2008). This suggests trout fisheries may be impacted by NZM infestation.

**Economic:** The economic costs of NZM infestations are primarily speculative and have received little quantitative



investigation. Due to their high densities, NZM may cause some degree of economic damage via biofouling structures such as water intake pipes; however, this has not commonly been reported and is likely minor in comparison to other invasive freshwater mollusks such as Zebra Mussels (Proctor et al. 2007).

Likely, the most significant economic cost of NZM invasions is related to fisheries because NZM are known to reduce the weight and quality of game fish species by food web disruption (Poirier 2015) and are of particular concern to trout fisheries (Vincent and Bakker 2008). NZM infestations can also impair fish hatchery function and increase operation costs (Oliver et al. 2021).

## Management

---

Management Goals: NZM presently have an expanding distribution in Pennsylvania. Therefore, the primary management goal must be to contain infestations.

### Containment and Prevention Actions:

- Coordinate with partners to conduct surveys/monitoring within invaded watersheds to determine the extent of NZM infestations, and initiate surveys of waters at-risk of infestation.
- Continue public education efforts to acquaint anglers with the threat of, and measures to prevent, the spread of NZM.
- Routinely monitor fish hatcheries at-risk of NZM contamination due to the presence of this species in nearby waters. Continue following established

Pennsylvania Fish and Boat Commission (PFBC) biosecurity protocols and response plans for fish rearing and stocking and incident response plans as needed.

- Support or initiate research in Pennsylvania further evaluating the impacts of NZM on native species and coldwater fisheries, and the habitat preferences of this species in order to predict areas of high risk for invasion.
- Continue installing NZM-specific signage developed by the PFBC and PA Sea Grant at locations infested by NZM and consider the installation of wader cleaning stations at infested sites. Established gear disinfection protocols include the following: freezing gear for a minimum of six hours, soaking gear in hot (>120°F) for at least five minutes, or completely drying gear for at least five days.
- Encourage the incident reporting of aquatic invasive/nuisance species such as NZM within Pennsylvania. Online reporting can now be conducted at the following PFBC web site: <https://www.fishandboat.com/Conservation/AIS/Pages/default.aspx> as well as PA iMapInvasives at: <https://www.paimapinvasives.org/> and at the national level, USGS Nonindigenous Aquatic Species website: <https://nas.er.usgs.gov/SightingReport.aspx>
- Keep informed with research concerning the use of biological/chemical controls (see Rapid Response Options below).





Rapid Response Options:

- Several physical or chemical treatment options have been evaluated for killing or preventing the dispersal of NZM within fish hatcheries; however, these are largely ineffective against NZM control due to the ability of this species to seal its shell during adverse chemical conditions (Oplinger et al. 2009). Copper sulfate-based treatments were recently demonstrated to be effective at eradicating NZM but required long-term (weeks) of treatment to be effective within hatchery raceways (Oliver et al. 2021). No effective methods of control appear to have been demonstrated for streams, rivers, large ponds, or lakes.

---

## References

---

- Benson, A.J., Kipp, R.M., Larson, J. and Fusaro, A. 2023. *Potamopyrgus antipodarum* (J.E. Gray, 1853): U.S. Geological Survey, Nonindigenous Aquatic Species Database, Gainesville, FL.  
<https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=1008>. Accessed 11/2/2023.
- Bilka, R.H. and Levri, E.P. 2013. The invasive New Zealand Mudsnail (*Potamopyrgus antipodarum*) grows faster when consuming periphyton compared to detritus. *Journal of the Pennsylvania Academy of Science*. 87: 125-128.
- Cada, C.A. 2004. Interactions between the invasive New Zealand Mudsnail, *Potamopyrgus antipodarum*, Baetid mayflies, and fish predators. Master

of Science Thesis, Montana State University. 126 pp.

- Hartzell, S.M. and Frederick, J.R. 2023. First Records of the Invasive New Zealand Mudsnail (*Potamopyrgus antipodarum*) in the Potomac River Basin. *Northeastern Naturalist*. 30: N13-N16.
- Hartzell, S.M. and Macelko, N. In Press. Range Expansion of the Invasive New Zealand Mudsnail (*Potamopyrgus antipodarum*) in the Susquehanna and Delaware River Basins of Pennsylvania. *Journal of the Pennsylvania Academy of Science*.
- Karens, B.L., Dybdahl, M.F., Gangloff, M.M., and Jannot, J.E. 2005. *Potamopyrgus antipodarum*: distribution, density, and effects on native macroinvertebrate assemblages in the Greater Yellowstone Ecosystem. *Journal of the North American Benthological Society*. 24: 123-138.
- Karens, B.L., Cada, C.A., and Zichovich, J. 2010. Asymmetrical behavioral interactions between the New Zealand Mudsnail, *Potamopyrgus antipodarum*, and scraping, collector-gathering, and collector-filtering macroinvertebrates. *Journal of Freshwater Ecology*. 25: 657-666.
- Krist, A.C. and Charles, C.C. 2012. The invasive New Zealand Mudsnail, *Potamopyrgus antipodarum*, is an effective grazer of algae and altered the assemblage of diatoms more than



- native grazers. *Hydrobiologia*. 694: 143-151.
- Larson, M.D. and Black, A.R. 2016. Assessing interactions among native snails and the invasive New Zealand Mud Snail, *Potamopyrgus antipodarum*, using grazing experiments and stable isotope analysis. *Hydrobiologia*. 763: 147-159.
- Levri, E.P., Kelly, A.A., and Love, E. 2007. The invasive New Zealand Mud Snail (*Potamopyrgus antipodarum*) in Lake Erie. *Journal of Great Lakes Research* 33: 1-6.
- Levri, E.P. Macelko, N., Brindle, B., Levri, J.E., Dolney, T.J., and Li, X. 2020. The invasive New Zealand Mud Snail *Potamopyrgus antipodarum* (J.E. Gray, 1843) in central Pennsylvania. *BioInvasions Records*. 9: 109-119.
- Oliver, D.C., Loubere, A.D., and Sorenson, J.A. 2021. Efficacy of low-dose EarthTec® QZ treatment for the control of New Zealand mud snails *Potamopyrgus antipodarum* in a hatchery environment. *Management of Biological Invasions*. 12: 85-95.
- Oplinger, R.W., Brown, P., and Wagner, E.J. 2009. Effect of sodium chloride, tricaine methanesulfonate, and light on New Zealand Mud Snail behavior, survival of snails defecated from rainbow trout, and effects of epsom salt on snail elimination rate. *North American Journal of Aquaculture*. 71: 151-164.
- Pourier, J. 2015. New Zealand Mudsnail surveys at Lower Columbia River Basin National Fish Hatcheries. U.S. Fish and Wildlife Service Columbia River Fisheries Program. 39 pp.
- Proctor, T., et al. (eds). 2007. *National management and control plan for the New Zealand Mudsnail (Potamopyrgus antipodarum)*. Aquatic Nuisance Species Task Force New Zealand Mudsnail Management and Control Plan Working Group. 100 pp.
- Richards, D.C., O'Connell, P., and Cazier Shinn, D. 2004. Simple control method to limit the spread of the New Zealand Mudsnail, *Potamopyrgus antipodarum*. *North American Journal of Fisheries Management*. 24: 114-117.
- Vinson, M.R. and Baker, M.A. 2008. Poor growth of rainbow trout fed New Zealand Mudsnails *Potamopyrgus antipodarum*. *North American Journal of Fisheries Management* 28: 701-709.
-